

Massachusetts Deer Browse Impact Survey

Property Name: **Bolton Flats and small section of Oxbow** Town(s): **Harvard**

Ownership: **DFG and USFWS**

General Surroundings: Rural/Forest/Ag **Low-residential** Med-residential High-residential

General level of use by public: Low **Moderate** Heavy Unknown

Is this property open to hunting? **Yes** No Unknown Special Hunting Restrictions: **Archery by permit only on Oxbow**

General level of nearby hunting? None **Limited** Moderate Heavy Unknown

Date of site visit **8/17/17**

Name(s) conducting survey: **Stainbrook**

Description of the forest on the property: (e.g., conifer, mixed conifer/ hardwood, hardwood dominated; what are the dominant trees in the forest canopy; is it closed or open, is the forest older- or younger-aged, is there active logging, etc):

Very swampy with few dry areas to survey with tree cover. Mostly aspen in dry areas. A lot of invasive plants like oriental bittersweet and glossy buckthorn.

Survey Instructions: Using the worksheet on the back, walk through a representative area (10-50%) of the forest to come up with an average level of impact on a property. Avoid areas near trails and roads as these are not representative. Pay special attention to what is able to grow and which tree species are being browsed, especially in canopy gaps (e.g., areas where the sun is able to reach the forest floor from fallen trees, recent cuts, etc.). Take a few pictures showing overall forest floor as well as specific cases of browsing, if present. Please also record a GPS track or attach a map of the property with areas surveyed marked. After filling out the survey on the back, check the box below that best describes the general level of forest impacts from deer browsing. Please email a copy of the survey, GPS track/map, and pictures to david.stainbrook@state.ma.us.

General level of forest impacts from deer browsing on the property (*check one*):

☐ **1: Little to No Impact**

Preferred hardwood tree seedlings and saplings such as maple, oak, ash, and hickory are growing up to and above 6 feet tall, with little to no sign of browsing. The shrub and herbaceous layers are well developed and show little to no impact (e.g., Pink Lady's Slipper, Trilliums, and Wild Sarsaparilla are present)

☐ **2: Between Little to No Impact and Moderately Impacted**

☐ **3: Moderately Impacted**

Preferred hardwood tree seedlings and saplings such as maple, oak, ash, and hickory are growing up to and above 6 feet tall, but most do show some signs of browsing. Some moderately preferred trees (e.g., cherry) may show slight browsing, but no sign of browsing on low-preference tree species such as American beech and white pine. Some preferred to moderately preferred shrubs show evidence of browsing (e.g., viburnums) and preferred herbaceous plants (e.g., Trilliums, Pink Lady's Slipper, Canada Mayflower, and Wild Sarsaparilla) are present, but show some signs of flowering parts removed.

☐ **4: Between Moderately Impacted and Impacted**

☐ **5: Impacted**

Preferred hardwood tree seedlings and saplings such as maple, oak, and ash are not common, and when present, show signs of moderate to heavy browsing. In their place are other tree species such as American beech, white pine, cherry, birch, etc., which may show evidence of browsing. Some low-preference shrubs show evidence of browsing (e.g., blueberry, Glossy Buckthorn). Few preferred shrubs may be found scattered in the understory and Hay-scented Fern, low-bush blueberry, huckleberry, grasses/sedges, and barberry may dominate large sections of the forest floor.

☐ **6: Between Impacted and Heavily Impacted**

☐ **7: Heavily Impacted**

Tree seedlings and saplings preferred by deer are almost non-existent, and when present show signs of heavy browsing. Less preferred shrubs and trees show signs of browsing and/or most saplings are unable to grow above 6ft. Low preference trees, such as white pine may show evidence of browsing. A browse line is often visible below 6ft. Foliage of native shrubs and wildflowers are very limited. Ferns, grasses/sedges, and non-preferred invasive plants such as barberry may dominate the forest floor.

Classify average level of browsing on seedlings/saplings for the following species (minimum 20 stems):


Preferred Species:

| | | | | | | | |
|-----------------|---|---|---|---|---|---|------------------|
| Oak | 1 | 2 | 3 | 4 | 5 | Present in canopy, few in understory | Not Present/Seen |
| Red/Sugar Maple | 1 | 2 | 3 | 4 | 5 | Present in canopy, few in understory | Not Present/Seen |

Less Preferred Species:

| | | | | | | | |
|---------------------|---|---|---|---|---|---|------------------|
| Eastern Hemlock | 1 | 2 | 3 | 4 | 5 | Present in canopy, few in understory | Not Present/Seen |
| Birch | 1 | 2 | 3 | 4 | 5 | Present in canopy, few in understory | Not Present/Seen |
| American Beech | 1 | 2 | 3 | 4 | 5 | Present in canopy, few in understory | Not Present/Seen |
| Black Cherry | 1 | 2 | 3 | 4 | 5 | Present in canopy, few in understory | Not Present/Seen |
| White Pine | 1 | 2 | 3 | 4 | 5 | Present in canopy, few in understory | Not Present/Seen |
| Other: <u>Aspen</u> | 1 | 2 | 3 | 4 | 5 | Present in canopy, few in understory | Not Present/Seen |

Browsing Score




General level of deer sign on the property (e.g., scat, tracks, etc.): Low Medium High Unknown

Is there any evidence of deer impacts in surrounding areas (e.g., homes with landscaping damage)?

None Very Limited Moderate Heavy Unknown

Additional Notes (e.g., describe herbaceous layer):

Very sparse with some Canada mayflower and wild sarsaparilla. Some browsing on ferns and glossy buckthorn, which were quite abundant. There appears to be difficult hunting access to the area.





Overall look of the forest showed forest heavily dominated by glossy buckthorn and other invasive plants, like oriental bittersweet and Japanese barberry.



Browsing on glossy buckthorn, typically not preferred by deer.



Browsing on fern (species unknown).



Browsing on Oriental bitterweet, typically not preferred by deer.



Browsing on low-bush blueberry, typically not preferred by deer.



A few small white oak saplings, but they were a few years old and less than a foot tall.